

# Neurology<sup>®</sup>

## **MRI-based biomarkers of preclinical AD : An Alzheimer signature**

Susan M. Resnick and Philip Scheltens

*Neurology*; Published online before print December 21, 2011;  
DOI 10.1212/WNL.0b013e31824237a5

**This information is current as of December 22, 2011**

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://www.neurology.org/content/early/2011/12/21/WNL.0b013e31824237a5>

*Neurology*® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2011 by AAN Enterprises, Inc. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.



# MRI-based biomarkers of preclinical AD

## An Alzheimer signature

Susan M. Resnick, PhD  
Philip Scheltens, MD,  
PhD

Correspondence & reprint  
requests to Dr. Resnick:  
susan.resnick@nih.gov

*Neurology*® 2012;78:80–81

Recently proposed criteria for prodromal Alzheimer disease (AD),<sup>1</sup> mild cognitive impairment (MCI) due to AD,<sup>2</sup> and probable AD dementia<sup>3</sup> incorporate molecular evidence of amyloid- $\beta$  ( $A\beta$ ) pathology and also consider measures of brain structure and function as supportive biomarkers. MRI-based biomarkers are also proposed as supportive evidence for a diagnosis of preclinical AD<sup>4</sup> in cognitively normal individuals, posing new challenges for definition, validation, and interpretation of these biomarkers. A number of MRI-based biomarkers, such as volumes of hippocampus and entorhinal cortex, show robust differences between groups of patients with AD and controls. However, these measures have limited sensitivity and specificity in predicting who will develop AD in individual patients. The development and validation of new structural neuroimaging biomarkers sensitive to early changes in the disease process will be critical to implementation of the new research criteria for preclinical AD.

In this issue of *Neurology*®, Dickerson and Wolk<sup>5</sup> investigate the utility of a potential MRI biomarker of neurodegeneration, based on a previously identified set of 9 brain regions that show cortical thinning in AD, MCI, and cognitively normal individuals with  $A\beta$  deposition on PET imaging. Using MRI data from 159 cognitively normal individuals in the Alzheimer's Disease Neuroimaging Initiative (ADNI), the authors test the hypothesis that individuals at high risk for preclinical AD based on this "AD signature" of regional cortical thinning would be at increased risk for subsequent cognitive decline over a 3-year follow-up. Based on the AD signature scores, cognitively normal individuals were classified as high, average, or low risk of preclinical AD. Significant cognitive decline was operationalized as an increase of  $\geq 1.0$  on the Clinical Dementia Rating Scale Sum of Boxes score and a decline of  $\geq 1$  SD on one of the following measures: immediate recall on a measure of verbal learning, verbal delayed free recall, or executive function measured by Trails B. The au-

thors report that normal individuals at high risk for preclinical AD are more likely to show cognitive decline, with risks of cognitive decline equal to 21% in the high-risk, 7% in the average-risk, and 0% in the low-risk group. This represents a 3-fold increase in risk for cognitive decline for each 1 SD of cortical thinning. In addition, 60% of the high-risk group had CSF characteristics (low CSF  $A\beta$ ) suggestive of AD in the subsample for which CSF data were available.

Dickerson and Wolk provide additional support for the utility of MRI-based biomarkers in identifying cognitively normal individuals at increased risk for cognitive impairment and AD. Importantly, the regions contributing to the cortical thinning "AD signature" were defined in previous work by the authors and were applied in the current investigation to a new sample, the ADNI controls who had prospective cognitive follow-ups. The use of multiple brain regions to develop a composite measure of brain abnormality is an additional strength of this work. The authors show that their composite AD signature, compared with entorhinal cortex thickness, provides better discrimination between individuals who show cognitive decline vs stable performance at follow-up. Composite measures of networks of brain regions that discriminate between normal and pathologic brains are likely to yield more informative prediction on an individual patient basis as compared to single regions alone, as shown in prior studies using support vector machine learning methods to distinguish spatial patterns of abnormality that identify the earliest stages of AD.<sup>6</sup>

Although the sample size in the current study was small (only 14 individuals with complete cognitive follow-up data met the criteria for high risk for preclinical AD) and longer follow-up intervals will be required to validate the AD signature against other (imaging) biomarkers, the ability to identify cognitively normal individuals at higher risk for subsequent cognitive decline is an important step toward

See page 84

From the National Institute on Aging (S.M.R.), Baltimore, MD; and Alzheimer Center (P.S.), Department of Neurology, VU University Medical Center, Amsterdam, the Netherlands.

*Disclosure:* Author disclosures are provided at the end of the editorial.

implementing and evaluating the new criteria for preclinical AD. Combinations of A $\beta$  biomarkers and structural imaging biomarkers of subtle regional neurodegeneration, along with their time course, may also help distinguish cognitively normal individuals with A $\beta$  pathology who will develop disease from those who will remain asymptomatic during their lifetime. Data from autopsy and PET amyloid imaging studies show that about 30% of individuals with normal cognitive function have substantial amyloid pathology. Although some argue that this group represents preclinical AD,<sup>2</sup> the absence of antemortem cognitive decline in prospective autopsy samples of “asymptomatic AD” suggests that a subgroup of individuals may demonstrate cognitive resilience despite substantial pathology.<sup>7</sup> In this group in particular, the presence of the “AD signature” in the structural, downstream, imaging marker may be predictive of who will actually decline. Distinguishing individuals who will decline from those who will remain asymptomatic is essential for patient selection and monitoring in therapeutic trials and for identification of individuals most likely to benefit from potential treatments to delay the onset of cognitive impairment.

#### AUTHOR CONTRIBUTIONS

Dr. Resnick: drafting/revising the manuscript. Dr. Scheltens: drafting/revising the manuscript.

#### DISCLOSURE

Dr. Resnick serves as Action Editor for *Brain and Cognition*; receives research support from the NIH/NIA Intramural Research Program; her spouse serves on a scientific advisory board and as a consultant for Amgen; and receives research support from Eli Lilly and Company, Roche, Amgen, Avid Radiopharmaceuticals, Inc., Johnson & Johnson, Lundbeck

Inc., Synosia Therapeutics, GE Healthcare, and the NIH. Dr. Scheltens serves on scientific advisory boards for Danone Research, Wyeth/Elan Corporation, Bristol-Myers Squibb, Genentech, Inc., Pfizer Inc, GE Healthcare, and Janssen AI; has received speaker honoraria from Lundbeck Inc. and Nutricia; serves as Book Review Editor for *Alzheimer's Disease and Associated Disorders* and on the editorial board of *Dementia Geriatric Cognitive Disorders*; serves as a consultant for Pfizer Inc, GE Healthcare, Avid Radiopharmaceuticals, Inc./Eli Lilly and Company, Genentech, Inc., and Janssen AI; and receives research support from Alzheimer Nederland and Stichting VUmc fonds.

#### REFERENCES

1. Dubois B, Feldman HH, Jacova C, et al. Revising the definition of Alzheimer's disease: a new lexicon. *Lancet Neurol* 2010;9:1118–1127.
2. Albert MS, DeKosky ST, Dickson D, et al. The diagnosis of mild cognitive impairment due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement* 2011;7:270–279.
3. McKhann GM, Knopman DS, Chertkow H, et al. The diagnosis of dementia due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement* 2011;7:263–269.
4. Sperling RA, Aisen PS, Beckett LA, et al. Toward defining the preclinical stages of Alzheimer's disease: recommendations from the National Institute on Aging Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement* 2011;7:280–292.
5. Dickerson BC, Wolk DA. MRI cortical thickness biomarker predicts AD-like CSF and cognitive decline in normal adults. *Neurology* 2012;78:84–90.
6. Davatzikos C, Fan Y, Wu X, Shen D, Resnick SM. Pattern classification of MRI detects prodromal Alzheimer's disease. *Neurobiol Aging* 2008;29:514–523.
7. Resnick SM, Sojkova J. Amyloid imaging and memory change for prediction of cognitive impairment. *Alzheimers Res Ther* 2011;3:3.

## MRI-based biomarkers of preclinical AD : An Alzheimer signature

Susan M. Resnick and Philip Scheltens

*Neurology*; Published online before print December 21, 2011;

DOI 10.1212/WNL.0b013e31824237a5

**This information is current as of December 22, 2011**

<b>Updated Information &amp; Services</b>	including high resolution figures, can be found at: <a href="http://www.neurology.org/content/early/2011/12/21/WNL.0b013e31824237a5">http://www.neurology.org/content/early/2011/12/21/WNL.0b013e31824237a5</a>
<b>Supplementary Material</b>	Supplementary material can be found at: <a href="http://www.neurology.org/content/suppl/2011/12/21/WNL.0b013e31824237a5.DC1.html">http://www.neurology.org/content/suppl/2011/12/21/WNL.0b013e31824237a5.DC1.html</a>
<b>Permissions &amp; Licensing</b>	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: <a href="http://www.neurology.org/misc/about.xhtml#permissions">http://www.neurology.org/misc/about.xhtml#permissions</a>
<b>Reprints</b>	Information about ordering reprints can be found online: <a href="http://www.neurology.org/misc/addir.xhtml#reprintsus">http://www.neurology.org/misc/addir.xhtml#reprintsus</a>

